

# Pion-LT/Kaon-LT Collaboration Meeting

**Muhammad Junaid**  
**Ph.D. Student**  
**Department of Physics**  
**University of Regina, Canada**

# LTSep Pre-Analysis

- ❑ Working on physics setting:  $Q2 = 3.85$ ,  $W = 2.62$ ,  $t = 0.21$  (2 epsilons)
- ❑ Finalized the following studies before the LTSep analysis:
  - Missing mass offset and cut determination
  - Diamond cut determination
  - t-resolution check
  - t-binning
  - phi-binning
  - Data yields
  - SIMC yields
  - Data/SIMC comparison and ratios
  - Average kinematics and ratios calculation

# LTSep Analysis

❑ Next steps are listed as follows:

- Unseparated cross-section calculations
- Model iterations
- Rosenbluth equation fitting
- L/T separated cross-section calculations
- Pion Form Factor measurements

# LTSep Analysis

- ❑ Calculated kinematic variables for each t-bin for both data and SIMC.
  - Error weighted average of Q2 for high and low epsilon (combined – low and epsilon settings)
  - Error weighted average of W for high and low epsilon (combined – low and epsilon settings)
  - Error weighted average of epsilon for high and low epsilon (combined – low and epsilon settings)
  - Error weighted average of theta for high and low epsilon (combined – low and epsilon settings)
  - Average t-central for high and low epsilon (combined – low and epsilon settings)
  - Average phi-central for high and low epsilon (combined – low and epsilon settings)
- ❑ Calculated ratios (DATA/SIMC) for each t & phi-bin setting-by-setting, separately.
- ❑ Calculated error weighted average of ratios for t & phi-bin for both high and low epsilon (combined - center, right, and left settings)
- ❑ Error-weighted average calculations:


$$w_i = \frac{1}{e_i^2}$$
$$\bar{x} = \frac{\sum (x_i \cdot w_i)}{\sum w_i}$$
$$\sigma_{\bar{x}} = \sqrt{\frac{1}{\sum w_i}}$$

# Iteration 0

# Physics Ratios

No iterations


Physics Setting:  
Q3p85\_W2p62\_t0p21\_loweps



# Physics Ratios

No iterations


Physics Setting:  
Q3p85\_W2p62\_t0p21\_higheps



# Un-separated Cross-sections

No iterations

Physics Setting:  
Q3p85\_W2p62\_t0p21  
Iteration: 0






# LT-separated Cross-sections


No iterations

Physics Setting:  
Q3p85\_W2p62\_t0p21  
Iteration: 0



# LT-separated Cross-sections

No iterations



# LTSep Analysis

- ❑ Working on physics setting: “ $Q^2 = 3.85$ ,  $W = 2.62$ ,  $t = 0.21$  (2 epsilons)”
- ❑ The following studies have been finalized for Pion Form Factor measurement:
  - Unseparated cross-section calculations
  - Model iterations
  - Rosenbluth equation fitting
  - L/T separated cross-section calculations
- ❑ In progress:
  - ❖ Working on iterations scripts